arXiv:nlin/0106032v1 [nlin.SI] 20 Jun 2001

Elliptic Solitons with Free Constants and their Isospectral Deformations

Yurii V. Brezhnev
Theoretical Physics Department,
Kaliningrad State University,
A. Nevsky st. 14, 236041, Kaliningrad, Russia
brezhnev@mail.ru

*

February 8, 2008

Abstract

New exact solvable elliptic potentials with free constants for the spectral problems of the third order are found. A time dependence of such potentials gives their isospectral deformations and solutions of nonlinear integrable equations.

Key words: elliptic solitons, finite-gap potentials, algebraic curves, spectral problems.

1 Introduction

In 1987, Verdier [1, 2] found new finite-gap potential for the Schrödinger equation

$$\Psi'' - u(x)\,\Psi = \lambda\,\Psi \tag{1}$$

in elliptic functions $u(x) = 6\,\wp(x) + 2\,\wp(x-\omega_i)$, differed from the well known Lamé potentials $u = n(n+1)\,\wp(x)$ and their isospectral deformations. Soon, the wide classes of new elliptic solitons (in terminology of Treibich and Verdier) were found. Later, three main approachs in the theory of elliptic solitons have been developed, including the scalar and matrix spectral problems. 1) The classical approach of Hermite–Halphen, based on the ansatz for the Ψ-function [3–7]; 2) An analysis of the algebraic curves of the special form — curves being a covering over tori [8, 18, 14]; 3) The methods, touching the Picard's theorem and hierarchies of the higher stationary nonlinear differential equations [9–13].

---

*Supported by Russian Basic Research Foundation with Grant 00–01–00782

There are the following main problems in the theory of elliptic solitons: the search for the potentials $u_i(x)$, formula for the Ψ-function, corresponding algebraic curve and its realization in the form of covering of torus, isospectral deformations of u_i with respect to integrable nonlinear equations.

Originally (1872), Hermite developed an approach to the integration of the Lamé potentials for (1) using an ansatz for the Ψ-function as a meromorphic two-periodic function of the second kind by Picard [3, 26] (Hermite used Jacobi's H, Θ, Z-functions):

$$\Psi = \frac{\sigma(x-\alpha_1)\cdots\sigma(x-\alpha_m)}{\sigma(x-\Omega_1)\cdots\sigma(x-\Omega_m)}\, e^{k(\lambda)x}, \tag{2}$$

where the poles Ω_j of the potential $u_i(x)$ independent of λ. In such ansatz, zeroes of the Ψ-function α_i depend on the spectral parameter λ and are determined from the *several* essentially *transcendental* equations. By this reason the ansatz (2) is not convenient and not effective. Later, Hermite and Halphen considered the following ansatz

$$\Psi = e^{kx} \sum_{n,\,j} a_{nj}\, \Phi^{(n)}(x-\Omega_j;\alpha), \tag{3}$$

where the Φ-function (*élément simple* by Halphen) is defined by the expression

$$\Phi(x;\alpha) = \frac{\sigma(\alpha-x)}{\sigma(\alpha)\,\sigma(x)}\, e^{\zeta(\alpha)x}.$$

Thus, instead of unknown parameters $\alpha_i(\lambda)$, $k(\lambda)$ in (2) we have only two ones k, α in the framework of ansatz (3). Indeed, after the substitution (3) into the spectral problem

$$\hat{L}\Psi \equiv \frac{d^n}{dx^n}\Psi(x;\lambda) + u_1(x,\lambda)\,\frac{d^{n-1}}{dx^{n-1}}\Psi(x;\lambda) + \cdots + u_n(x,\lambda)\,\Psi(x;\lambda) = 0 \tag{4}$$

and expanding the principal part of poles at $x = \Omega_j$, we will have equations being polynomial in k and transcendental in α. In fact, α is a parameter of the covered torus

$$\wp'(\alpha)^2 = 4\,\wp(\alpha) - g_2\,\wp(\alpha) - g_3.$$

Evidently, that the obtained equations are linear ones with respect to a_{nj} and therefore elementary solvable.

The classical potentials of Lamé, Halphen, Treibich–Verdier and others have a fixed location of poles. So, considering the potential

$$u(x) = 2\,\wp(x) + 2\,\wp(x-a) + 2\,\wp(x-b)$$

for the spectral problem (1), we get the restriction: $b = -a$, $\wp''(a) = 0$. In this paper, we find new potentials containing free parameters and poles (see also the most recent information in [25]). Assuming their dependence on time t we will obtain the isospectral deformation of initial condition $u(x)$ as a solution of the corresponding nonlinear integrable equation.

2 Algorithm for the solution of the spectral problems

The general algorithm for integrating the spectral problem (4) with elliptic coefficients u_i of x and arbitrary of λ was proposed by the author in [27]. It has a pure algebraic characterization and is reduced to a system of polynomial equations solvable by the polynomial algebra methods (Gröbner bases, characteristic sets and others [16, 17]). Indeed, the equations on poles of the expression $\hat{L}\Psi$, where the Ψ-function is given by the formula (3), contain the following unknown quantities:

$$k,\ \wp(\alpha),\ \wp'(\alpha),\ \wp(\Omega_i-\Omega_j),\ \wp'(\Omega_i-\Omega_j),\ \zeta(\alpha),\ \zeta(\Omega_i-\Omega_j),\ \Phi(\Omega_i-\Omega_j;\alpha).$$

Using the addition theorems for elliptic functions and two important equalities

$$\Phi(x+z;\alpha)=\frac{1}{2}\,\frac{\Phi(x;\alpha)\,\Phi(z;\alpha)}{\wp_x-\wp_z}\left(\frac{\wp'_\alpha+\wp'_x}{\wp_\alpha-\wp_x}-\frac{\wp'_\alpha+\wp'_z}{\wp_\alpha-\wp_z}\right),\quad \Phi(\Omega;\alpha)\,\Phi(-\Omega;\alpha)=\wp_\alpha-\wp_\Omega,$$

where we introduce the short notation $\wp_\alpha\equiv\wp(\alpha)$ etc., the above mentioned equations on poles are "algebraized": they are transformed into the polynomial ones with respect to the variables k, $\wp(\alpha)$, $\wp'(\alpha)$, $\wp(\Omega_i)$, $\wp'(\Omega_i)$ (see [27] for the details). Thus, we have a polynomial ideal in a ring $\mathbb{Q}(\lambda,g_2,g_3)[k,\wp_\alpha,\wp'_\alpha,\wp_\Omega,\wp'_\Omega]$ and the final solution of the spectral problem (4) is defined by the structure of this ideal [15, 17, 27].

3 Elliptic solitons with free constants

Let us consider the spectral problem

$$\Psi'''-u(x)\,\Psi'=\lambda\,\Psi. \tag{5}$$

Here, the potentials with only pole in a parallelogramm of periods have not unified form like the Lamé potentials. Indeed, the general 1-pole elliptic potential has the form

$$u(x)=A\,\wp(x)+B. \tag{6}$$

Investigating (6) in the framework of the ansatzs

$$\Psi=\Big(\Phi(x;\alpha)+a_1\,\Phi'(x;\alpha)+\cdots+a_n\,\Phi^{(n)}(x;\alpha)\Big)\,e^{kx} \tag{7}$$

under $n=0,1,2,3,4$ we get:

- $n=0$: $u=6\,\wp(x)+B$ [22, p. 509]. B is an arbitrary constant, genus $g=2$;
- $n=1$: $u=12\,\wp(x)$. The degenerated curve is of genus $g=1$;
- $n=2$: No solution. λ must be zero [7];
- $n=3$: $u=30\,\wp(x)\pm3\sqrt{3\,g_2}$. Genus $g=3$;

- $n = 4$: $u = 42\,\wp(x)$. Genus $g = 6$.

All the associated curves are the trigonal ones. λ is a meromorphic function with only 3-rd order pole. The 2-pole elliptic potentials in (5) have the form

$$u = A\,\wp(x) + B\,\wp(x-\Omega) + C\,\zeta(x) - C\,\zeta(x-\Omega) + D.$$

Using the Θ-functional approach, one can show that the finite-gap potential for (5) is a second logarithmic derivative of Θ and thus, must not has residues. Therefore $C = 0$. The simplest nontrivial example is

$$u = 6\,\wp(x) + 6\,\wp(x-\Omega). \tag{8}$$

The corresponding solution is as follows [27]:

$$\begin{gathered}\Psi(x) = \Big(a\,\Phi(x;2\,\alpha) + \Phi(\Omega;2\,\alpha)\,\Phi(x-\Omega;2\,\alpha)\Big)e^{kx},\\ a = \zeta(2\,\alpha+\Omega) - 2\,\zeta(\alpha) - \zeta(\Omega), \quad k = 2\,\zeta(\alpha) - \zeta(2\,\alpha), \quad \lambda = -4\,\wp'(\alpha).\end{gathered} \tag{9}$$

The generalization of Halphen's equation

$$\Psi''' - ((n^2-1)\,\wp(x) + h)\,\Psi' - \tfrac{n^2-1}{2}\,\wp'(x)\,\Psi = \lambda\,\Psi$$

was considered by Unterkofler [20] in the framework of the ansatz (2). The algebraic curves were written, but important formulas for the covering of torus $R(\wp(\alpha), \wp'(\alpha), \lambda) = 0$ (i.e. the transcendental equation for the determining of α by λ) have not been obtained.

Let us consider the spectral problem

$$\Psi''' - u(x)\,\Psi' - v(x)\Psi = \lambda\,\Psi. \tag{10}$$

The most general 1-pole elliptic potential for the equation (10) has the form:

$$\Psi''' - (a\,\wp(x) + d)\,\Psi' - (b\,\wp'(x) + c\,\wp(x))\,\Psi = \lambda\,\Psi. \tag{11}$$

The case $n = 0$ in the ansatz (7) yields the solution:

$$\Psi''' - 3\,(\wp(x) + c^2)\,\Psi' - \Big(\tfrac{3}{2}\,\wp'(x) + 3\,c\,\wp(x)\Big)\,\Psi = \lambda\,\Psi, \qquad \Psi = \Phi(x;\alpha)\,e^{cx}.$$

Under $n = 1$ in (7) we arrive at the nontrivial solution. Indeed, the first of three polynomials in the original base has the form:

$$(a-12)\,\Big((a-18)\,(k^2 - \wp_\alpha) + 2\,c\,k\Big) + 2\,(a-18)\,d + c^2 = 0, \tag{12}$$

where without loss of generality we set $b = 12 - a$. In general case, an attempt to solve the corresponding system of algebraic equations failed. For example, in some particular cases, all the solutions lead to the dependence of parameters a, c, d on spectral one λ.

But under $a = 12$ the polynomial (12) is simplified: the dependence on k, α disappeares. After the replacement $c \to 12\,c$, the determining system of equations takes the form

$$\begin{aligned}
&3\,k^4 - 4\,c\,k^3 - 6\,(3\,\wp + 2\,c^2)\,k^2 + 12\,(c\,\wp + \wp')\,k - 9\,\wp^2 + 12\,c^2\,\wp - 4\,c\,\wp' - 2\,\lambda\,c + 3\,g_2 = 0,\\
&4\,k^3 - 6\,c\,k^2 - 12\,(\wp + 2\,c^2)\,k + 6\,c\,\wp + 4\,\wp' + 24\,c^3 - \lambda = 0,\\
&\wp'^2 - 4\,\wp^3 + g_2\,\wp + g_3 = 0
\end{aligned} \tag{13}$$

and the solution for the remaining variables read as

$$\Psi''' - 12\,(\wp(x) + c^2)\,\Psi' - 12\,c\,\wp(x)\,\Psi = \lambda\,\Psi, \quad \Psi = \Big((k - 2\,c)\,\Phi(x;\alpha) + \Phi'(x;\alpha)\Big)e^{kx}. \tag{14}$$

The ideal (13) is not equal $\langle 1 \rangle$ in a ring $\mathbb{Q}(\lambda, c, g_2, g_3)[\wp, \wp', k]$. Thus, c is an arbitrary constant. For a completeness we give the expression for the algebraic curve $F(k,\lambda) = 0$ depended on c:

$$\begin{aligned}
&64\,(\lambda - 11\,c^3)(\lambda^2 + 32\,\lambda\,c^3 + 2^8\,c^6 - 108\,g_2\,c^2)\,\underline{k^3} - 48\,(c\,\lambda^3 + 4\,(3\,c^4 + g_2)\,\lambda^2 -\\
&4\,(96\,c^6 + 13\,g_2\,c^2 - 36\,g_3)\,c\,\lambda - 16\,(320\,c^8 - 127\,g_2\,c^4 + 99\,g_3\,c^2 + 9\,g_2^2)\,c^2)\,\underline{k^2} -\\
&48\,(11\,c\,\lambda^3 + 4\,(60\,c^4 + g_2)\,\lambda^2 - 4\,(192\,c^6 + 301\,g_2\,c^2 - 72\,g_3)\,c\,\lambda -\\
&16\,(1792\,c^8 - 676\,g_2\,c^4 + 198\,g_3\,c^2 - 9\,g_2^2)\,c^2)\,c\,\underline{k} - 27\,\lambda^4 - 352\,c^3\,\lambda^3 +\\
&24\,(160\,c^6 + 151\,g_2\,c^2 - 36\,g_3)\,\lambda^2 - 384\,(128\,c^8 + 112\,g_2\,c^4 - 18\,g_3\,c^2 + 3\,g_2^2)\,c\,\lambda -\\
&2^{17}11\,c^{12} + 2^{10}609\,g_2\,c^8 - 2^8 891\,g_3\,c^6 - 25776\,g_2^2\,c^4 + 2^7 3^5\,g_2\,g_3\,c^2 + 2^8\,(g_2^3 - 27\,g_3^2) = 0.
\end{aligned} \tag{15}$$

This curve has 8 finite points of ramification λ_i with indexes (2,1) and a ramification at infinity with index 3. Genus $g = 3$. The curve is not hyperelliptic one. The corresponding covering of torus $R(\wp(\alpha), \wp'(\alpha), \lambda) = 0$ is as follows:

$$\begin{aligned}
&32\,(\lambda - 11\,c^3)(\lambda^2 + 32\,\lambda\,c^3 + 2^8\,c^6 - 108\,g_2\,c^2)\,\underline{\wp'(\alpha)} - 48\,(c\,\lambda^3 + 4\,(3\,c^4 + g_2)\,\lambda^2 -\\
&4\,(96\,c^6 + 13\,g_2\,c^2 - 36\,g_3)\,c\,\lambda - 16\,(320\,c^8 - 127\,g_2\,c^4 + 99\,g_3\,c^2 + 9\,g_2^2)\,c^2)\,\underline{\wp(\alpha)} +\\
&\lambda^4 + 32\,c^3\,\lambda^3 - 24\,(g_2\,c^2 + 12\,g_3)\,\lambda^2 - 128\,(64\,c^8 + 6\,g_2\,c^4 - 9\,g_3\,c^2 + 3\,g_2^2)\,c\,\lambda -\\
&2^{16}\,c^{12} - 2^{11}3\,g_2\,c^8 - 94464\,g_3\,c^6 + 5520\,g_2^2\,c^4 + 2^7 81\,g_2\,g_3\,c^2 + 2^8\,(g_2^3 - 27\,g_3^2) = 0.
\end{aligned} \tag{16}$$

This equation as an elliptic function of α of the 3-rd order has three solutions $\alpha_{1,2,3}$ corresponding to three linear independent solutions of (14) with arbitrary λ, c. To all appearances, the potential (14) corresponds to the higher Boussinesq equations, but its complete finite-gap treatment is an open question. We revealed one more new case, when a polynomial system is simplified and a solution is not trivial. It is

$$\Psi''' - (18\,\wp(x) + d)\,\Psi' + 6\,\wp(x)\,\Psi = \lambda\,\Psi, \quad \Psi = \Big((3\,k^2 + 3\,\wp_\alpha - d)\,\Phi(x;\alpha) + 6\,k\,\Phi'(x;\alpha)\Big)e^{kx}.$$

The corresponding formulas for curve and covering type as (15) and (16) of genus $g = 4$ are too big that to display here. Another case

$$\psi''' - (6\,\wp(x) + d)\,\psi' - 6\,\wp'(x)\,\psi = \lambda\,\psi \tag{17}$$

is trivial in the sense that $\psi = \Psi'$ in (5) with $n = 0$ is the solution of (17).

The next nontrivial example is a simplest 2-pole potential in (10).

$$\Psi''' - 3\,(\wp(x) + \wp(x-\Omega) - \wp_{A})\,\Psi' - \frac{3}{2}\Big(\wp'(x) + \wp'(x-\Omega) + B\,\wp(x) - B\,\wp(x-\Omega)\Big)\,\Psi = \lambda\,\Psi. \tag{18}$$

This equation is integrable with arbitrary Ω and A (see [27, 23] for the solution) and the restriction $B^2 = 4\,\wp_{\Omega} - 4\,\wp_{A}$.

4 Isospectral deformations

If Ψ-function depends on a parameter t, then the arbitrariness of location of poles Ω_i in potential means its isospectral deformation with $\Omega_i(t)$. Let us consider examples.

The spectral problem (5) corresponds to the Sawada–Kotera equation [19]

$$u_t = u_{xxxxx} - 5\,u\,u_{xxx} - 5\,u_x\,u_{xx} + 5\,u^2\,u_x. \tag{19}$$

A deformation of the potential (8) is stationary. Chazy [21, p. 380] found the more general stationary solution of the equation (19) $u(x,t) = 6\,\wp_1 + 6\,\wp_2$, where we introduced the designations:

$$\wp_1 \equiv \wp(x - c\,t - \Omega;\, g_2, g_3), \quad \wp_2 \equiv \wp(x - c\,t - \tilde{\Omega};\, g_2, \tilde{g}_3), \quad c = -12\,g_2.$$

Now we use a connection between the equation (19) and the Kaup–Kupershmidt, Fordy–Gibbons [28] and Tzitzeica equations respectively

$$w_t = w_{xxxxx} - 5\,w\,w_{xxx} - \frac{25}{2}\,w_x w_{xx} + 5\,w^2\,w_x, \tag{20}$$

$$v_t = v_{xxxxx} - 5\,(v_x\,v_{xxx} + v^2\,v_{xxx} + v_{xx}^2 + v_x^3 + 4\,v\,v_x\,v_{xx} - v^4\,v_x), \tag{21}$$

$$\phi_{xt} = e^{\phi} - e^{-2\,\phi} \tag{22}$$

by means of the Miura maps

$$u = v^2 - v_x, \qquad w = v^2 + 2\,v_x, \qquad u = \phi_{xx} + \phi_x^2. \tag{23}$$

After the substitution $v = -\ln'\psi$, the first Miura's map in (23) is linearized and we arrive at a problem of integration of the Schrödinger equation without the spectral parameter:

$$\psi_{xx} = 6\,(\wp_1 + \wp_2)\,\psi.$$

Using the solution (9) under $\lambda = 0$ (the passage to the limit $\alpha \to \omega_i$) we will have

$$\Psi = C_1\,\Big(\zeta(x) - \zeta(x - \Omega)\Big) + C_2.$$

After the substitution $\Psi' = \psi$, we get the result (under the coinciding invariants $g_3 = \tilde{g}_3$):

$$\psi = \wp_1 - \wp_2.$$

The straightforward verification shows that it will be also valid under $g_3 \neq \tilde{g}_3$. Thus we obtain the 5-parametric stationary solutions of the equations (20, 21):

$$v(x,t) = \frac{\wp_1' - \wp_2'}{\wp_1 - \wp_2}, \qquad w(x,t) = -12\,(\wp_1 + \wp_2) + 3\left(\frac{\wp_1' - \wp_2'}{\wp_1 - \wp_2}\right)^2.$$

The same technique leads to the solution of the Tzitzeica equation (22), but the time dynamics is not stationary:

$$\phi(x,t) = \ln 2\,c + \ln\Big(\wp(x + c\,t - \Omega;\, g_2, g_3) - \wp(x - c\,t - \tilde{\Omega};\, g_2, \tilde{g}_3)\Big), \qquad 4\,(\tilde{g}_3 - g_3)\,c^3 = 1.$$

Let us consider the time deformation of 2-pole potential (18)

$$u(x,t) = 3\,\wp(x - \Omega_1) + 3\,\wp(x - \Omega_2) - 3\,\wp_A \tag{24}$$

with respect to the Boussinesq equation

$$3\,u_{tt} = \big(2\,u^2 - u_{xx}\big)_{xx}. \tag{25}$$

Substituting the ansatz (24) in (25) and equating the poles to zero we will obtain the equations on the poles $\Omega_{1,2}(t)$ — the Calogero–Moser system of two particles with a repulsion potential and immovable center of mass:

$$\left\{\begin{array}{rcl} \ddot{\Omega}_1 - 4\,\wp'(\Omega_1 - \Omega_2) &=& 0 \\ \ddot{\Omega}_2 + 4\,\wp'(\Omega_1 - \Omega_2) &=& 0, \end{array}\right. \qquad \left\{\begin{array}{rcl} \dot{\Omega}_1^2 - 4\,\wp(\Omega_1 - \Omega_2) + 4\,\wp_A &=& 0 \\ \dot{\Omega}_2^2 - 4\,\wp(\Omega_1 - \Omega_2) + 4\,\wp_A &=& 0. \end{array}\right. \tag{26}$$

Ruling out its trivial solution $\Omega_1 - \Omega_2 = C$, we get $\Omega_2 = C - \Omega_1$ and an equation on Ω_1:

$$\dot{\Omega}_1^2 = 4\,\wp(2\,\Omega_1 - C) - 4\,\wp_A. \tag{27}$$

The substitution $X = \wp(2\,\Omega_1 - C)$ transforms the equation (27) into the algebraic form:

$$\dot{X}^2 = 16\,(4\,X^3 - g_2\,X - g_3)(X - \wp_A). \tag{28}$$

(28) is the algebraic curve $F(X, \dot{X}) = 0$ of genus 1. After the birational change $X \to \tilde{X}$

$$X = \frac{3\,\wp_A\,\tilde{X} - 8\,g_2\,\wp_A - 12\,g_3}{3\,\tilde{X} - 48\,\wp_A^2 + 4\,g_2}$$

we obtain an equation on $\tilde{X}$ in the canonical Weierstrass's form

$$\left(\frac{d\tilde{X}}{dt}\right)^2 = 4\,\tilde{X}^3 - \tilde{g}_2\,\tilde{X} - \tilde{g}_3, \qquad \tilde{X}(t) = \wp(t - t_0;\, \tilde{g}_2,\, \tilde{g}_3) \equiv \tilde{\wp}(t - t_0)$$

with new invariants

$$\tilde{g}_2 = 2^8\,g_2\,\wp_A^2 + 768\,g_3\,\wp_A + \tfrac{64}{3}\,g_2^2,$$

$$\tilde{g}_3 = -2^{12}\, g_3\, \wp_A^3 - \frac{2^{11}}{3}\, g_2^2\, \wp_A^2 - 2^{10}\, g_2\, g_3\, \wp_A + \frac{2^9}{27}\, g_2^3 - 2^{10}\, g_3^2.$$

Thus, we get the complete solution of the equations (26):

$$\Omega_1(t) = \frac{1}{2}\, \wp^{-1}\!\left(\frac{3\, \wp_A\, \tilde{\wp}(t-t_0) - 8\, g_2\, \wp_A - 12\, g_3}{3\, \tilde{\wp}(t-t_0) - 48\, \wp_A^2 + 4\, g_2}\right) + \frac{C}{2}.$$

t_0, C are arbitrary constants and the elliptic integral $\wp^{-1}$ corresponds to the invariants g_2, g_3. See also [24] for the solution in the context of the solutions of the Kadomtsev–Petviashvili equation.

References

[1] J. L. Verdier: *New elliptic solitons*. Preprint, Paris, (1987).

[2] A. Treibich, J. L. Verdier: *Compt. Rend. Acad. Sci. Paris.* **311**, 51–54, (1990).

[3] E. T. Whittaker, G. N. Watson: *A Course of Modern Analysis*, Cambridge Univ. Press, Cambridge, 1986.

[4] C. Hermite: *Oeuvres de Charles Hermite* (Gautheir–Villar, Paris), vol. III, (1912).

[5] G. H. Halphen: *Mémoire sur la réduction des équations différentielles linéaires aux formes intégrables*, Mem. prés. l'Acad. Sci. France. **28**, 1–300 (1884).

[6] V. Z. Enol'skii, N. A. Kostov: *Acta Appl. Math.* **36**, 57–86 (1994).

[7] J. C. Eilbeck, V. Z. Enol'skii: *J. Math. Phys.* **35**(3), 1192–1201 (1994).

[8] A. O. Smirnov: *Acta Appl. Math.* **36**, 125–166 (1994).

[9] F. Gesztesy, R. Weikard: *Math. Nachr.* **176**, 73–91 (1995).

[10] F. Gesztesy, R. Weikard: *Compt. Rend. Acad. Sci. Paris.* **321**, 837–841 (1995).

[11] F. Gesztesy, R. Weikard: *Acta Math.* **176**, 73–107 (1996).

[12] F. Gesztesy, D. Race, R. Weikard: *J. London Math. Soc.* (2) **47**, 321–340 (1993).

[13] F. Gesztesy, R. Weikard: *Bull. Amer. Math. Soc.* **35**, 271–317 (1998).

[14] E. D. Belokolos, A. I. Bobenko, V. Z. Enol'skii, A. R. Its, V. B. Matveev: *Algebro–Geometric Approach to Nonlinear Integrable Equations*, Springer, Berlin 1994.

[15] Progress in Mathematics. **109**. *Computational Algebraic Geometry.* (F.Eyssette, A.Galligo, eds), Birkhäuser, 1993.

[16] B. Buchberger: *Gröbner Bases — an Algorithmic Method in Polynomial Ideal Theory*, in Multidimensional Systems Theory, N. K. Bose ed., D. Riedel, Dordrecht, 1985.

[17] A. M. Cohen, R. H. Cushman: *Gröbner Bases and Standard Monomial Theory.* Progress in Mathematics. **109**, 41–60 (1993).

[18] V. Z. Enol'skii, J. C. Eilbeck: *J. Phys. A: Math. Gen.* **28**, 1069–1088 (1995).

[19] K. Sawada, T. Kotera: *Progr. in Theor. Phys.* **51**(5), 1355–1367 (1974).

[20] K. Unterkofler: *Diff. Integral Eqs.* **14**(9), 2001

[21] J. Chazy: *Acta Math.* **34**, 317–385 (1911).

[22] Dr. E. Kamke: *Differentialgleichungen Lösunsmethoden und Lösungen*, New York, 1971.

[23] A. O. Smirnov: *Mat. Sbornik.* **190**(5), (1999) (in Russian).

[24] D. V. Chudnovsky: *J. Math. Phys.* **20**(12), 2416–2422 (1979).

[25] A. Treibich: *Beyond the exceptional covers and their canonical hyper-elliptic potentials.* Preprint 99–3, Univ. d'Artois, France, pp. 43 (1999).

[26] N. I. Akhiezer: *Elements of the Theory of Elliptic Functions*, Amer. Math. Soc., Providence, 1990.

[27] Y. V. Brezhnev: *Elliptic Solitons and Gröbner Bases*. http://xxx.lanl.gov, nlin.SI/0007028.

[28] A. P. Fordy, J. Gibbons: *Phys. Lett. A.* **75**, 325 (1980).